



## RESEARCH ARTICLE

**Antistress and antioxidant effects of *Borago officinalis* leaves**G. Mrudula<sup>1</sup>, B.M Vrushabendra Swamy<sup>2</sup>, K.N Jayaveera<sup>3</sup>, N.Anitha<sup>4</sup><sup>1</sup>Department of Pharmacology, Krupanidhi College of Pharmacy, Bangalore, Karnataka, India<sup>2</sup>Department of Pharmacology, East Point College of Pharmacy, Bangalore, Karnataka, India<sup>3</sup>Department of Chemistry, Jawaharlal Nehru Technological University, Anantapur, Andhra Pradesh, India<sup>4</sup>Department of Pharmacology, Sultan - Ul - Uloom College of Pharmacy, Hyderabad, Andhra Pradesh, India

Received 10 June 2014; Accepted 25 June 2014

**ABSTRACT**

The ethanolic extract of *Borago officinalis* (BO) (10, 25, 50 mg/kg, p.o) was tested for its Adaptogenic activity by using forced swimming test and *invitro* antioxidant activity by using 1, 1- Diphenyl, 2-Picryl – hydrazyl free radical, Nitric oxide scavenging activity and reducing power method. In antistress activity, pretreatment with BO extract significantly reduced the immobility time at 50 mg/kg was comparable to that of imipramine (20 mg/kg, i.p) treated group. Under *invitro* 1, 1- Diphenyl, 2-Picryl – hydrazyl free radical and nitric oxide free radicals are considerably inhibited in a dose dependent manner and there is increase in absorbing power indicates increase in reducing power. The results suggest that it could be used for the treatment of oxidative stress induced disorder.

**Key words:** *Borago officinalis*, Adaptogenic, Antistress, Antioxidant**INTRODUCTION:**

The word adaptogen is used by herbalists to refer to a natural herb product that increases the body's resistance to stresses such as trauma, anxiety and bodily fatigue. An adaptogen is nontoxic to the recipient. It produces a nonspecific response in the body - an increase in the power of resistance against multiple stressors including physical, chemical, or biological agents. It has a normalizing influence on physiology, irrespective of the direction of change from physiological norms caused by the stressor. Adaptogens have a normalizing effect on the body and are capable of either toning down the activity of hyper functioning systems or strengthening the activity of hypo functioning systems<sup>[1, 2]</sup>. Free radicals induced by per oxidation have gained much importance because of their use in several pathological conditions such as atherosclerosis, ischemia, liver disorder, neural disorder, metal toxicity and pesticide toxicity.

Together with other derivatives of oxygen, they are inevitable byproducts of biological redox reactions. Antioxidants may offer resistance against the oxidative stress by scavenging free radicals, inhibiting the lipid per oxidation and by other mechanisms and thus prevent diseases<sup>[3]</sup>.

The plant *Borago officinalis* (*B.officinalis* L.) family Boraginaceae, also known as "Star flower" is an annual herb originating in Syria, but naturalized throughout the Mediterranean region, as well as Asia, Europe, North

Africa and South America. In unani system of medicine the plant is very popular as Gaozaban. Naturopathic practitioners use borage for regulation of metabolism and the hormonal system<sup>[4]</sup>. In olden days it is used as nerve tonic as a treatment for depression, anxiety and grief. Leaves have been used as an adrenal tonic to counter the effects of steroid therapy. It also restores vitality during convalescence from illness<sup>[5, 6]</sup>. It was reported that ethanolic extract of defatted *B.officinalis* seeds possess phenolic acids, antioxidant and free radical scavenging activity<sup>[7, 8]</sup>. The crude extracts of borage possess anti spasmodic, bronchodilator, vasodilator and cardiac depressant activity<sup>[9]</sup>.

**METHODS:****Preparation of the extract:**

The leaves of *Borago officinalis* (BO) were purchased from Munnalal Dawasaz (Mfg of ayurvedic medicine) and shade dried. The plant was authenticated by an expert. The coarse powder was then macerated for 48 hrs and filtered using muslin cloth. The extracts were concentrated and evaporated to dryness under reduced pressure at 50°C.

**Animals:**

Albino rats (160-180 g) of either sex were used. They were kept in standard plastic animal cages in groups of 6-8 animals, with 12 hr of light and dark cycle in the

institutional animal house. The animals were fed with standard rodent diet and provided water *ad libitum*.

After one week of acclimatization the animals were used for further experiments. Approval from the institutional animal ethical committee for the usage of animals in the experiments was obtained as per the Indian CPCSEA guidelines. Approval number of institutional Animal Ethical Committee for this study was IEAC/SUCP/06/2007.

#### **Acute toxicity test:**

Acute toxicity of the preparation was determined using different groups of rats. The animals were fasted for 3 hrs prior to the experiment according to the recommended procedure (OECD guide no.425) [10]. As per guidelines, the animals were observed for 48 hrs for any mortality was observed and the preparation was found to be safe up to dose of 2000 mg/kg b.wt.

#### **Anti stress activity** [11,12,13]:

The rats of either sex 120-150 gm were divided into 5 groups. Each group containing six animals. Group I – Control vehicle 10 ml/kg b.wt p.o, Group II – Imipramine 20 mg/kg b.wt i.p, Group III – BO 10 mg/kg b.wt.p.o, Group IV – BO 25 mg/kg b.wt .p.o and Group V – BO 50 mg/kg b.wt .p.o. The BO extract was solubilised in 1% CMC and administered to the Groups (III, IV and V) animals for 15 days.

#### **Forced Swimming test** [14]:

The rats are individually forced to swim inside a vertical plexiglas's cylinder (height: 40cm; diameter: 18 cm, containing 15 cm of water maintained at 25°C). Rats are placed in the cylinders for the first time are initially highly active, vigorously swimming in circles, trying to climb the wall or diving to the bottom. After 2-3 min activity begins to subside and to be interspersed with phases of immobility or floating of increasing length. After 5-6 min immobility reaches a plateau where the rats remain immobile for approximately 80% of the time. After 15 min in the water the rats are removed and allowed to dry in a heated enclosure 32°C before being returned to the cages. They are again placed in the cylinder 24 h later and the total duration of immobility is measured during a 5 min test. Floating behavior during this 5min period has been found to be reproducible in different groups of rats. An animal is judged to be immobile whenever it remains floating passively in the water in a slightly hunched but upright position, its nose just above the surface. Standard drug Imipramine was administered one hour prior to testing. The results are shown in **Table 1** and depicted in **Figure 1**. Statistical Analysis Statistical analysis was performed using One way ANOVA followed by Dunnett's test. P values <0.05 were considered significant.

#### **InVitro Antioxidant Activity:**

##### **DPPH Radical Scavenging Activity** [15, 16, 17]:

DPPH is a stable free radical at room temperature and accepts an electron or hydrogen radical to become a stable diagnostic molecule. The reduction capacity of DPPH radical was determined by the decrease in its absorbance at 517 nm, which is induced by antioxidants. 0.1Mm solution of DPPH in ethanol was prepared and 0.1 ml of this solution was added to 3.0 ml of BO extract solution in water at different concentrations (10-100 ug/ml). Thirty minutes later, the absorbance was measured at 517 nm. Lower absorbance of the reaction mixture indicates higher free radical scavenging activity. Experimental was done in triplicate. The difference in the absorbance between the test and the control (DPPH in ethanol) was calculated and expressed as percent scavenging of DPPH radical.

The capability to scavenge the DPPH radical was calculated by using the following equation. Ascorbic acid was used as standard.

Scavenging effect (%) =  $[1 - \text{Abs. of sample} / \text{Abs. of control}] \times 100$

The results are shown in **Table 2** and depicted in **figure 2**

##### **Nitric oxide scavenging** [15, 16, 17]:

The nitric oxide generated from sodium nitroprusside in aqueous solution at physiological pH is inhibited by antioxidants which compete with oxygen to react with nitric oxide. The absorbance of the chromophore formed during diazotization of nitrite with sulphanilamide and subsequent coupling with naphthylethylenediamine (NED) was read at 546 nm. Nitric oxide was generated from sodium nitroprusside and measured by the Griess reagent. Sodium nitroprusside (5 mM) in phosphate buffer (pH 7.4) saline was mixed with different concentrations (10, 20, 40, 50, 60, 80, 100 ug/ml) of BO extract and incubated at 25°C for 150 min. To 1 ml of the incubated solution, 1 ml of Griess reagent (1% sulphanilamide, 2% O – Phosphoric acid and 0.1% naphthyl ethylene diamine dihydrochloride) was added. Absorbance was measured at 546 nm. % inhibition of OD was calculated by using the formula. Experiment was done in triplicate. The values are compared to standard drug Ascorbic acid. Capability to scavenge the nitric oxide radical was calculated by using the equation: % inhibition =  $[1 - \text{Abs. of sample} / \text{Abs. of control}] \times 100$

The results are shown in **Table 3** and depicted in **figure 3**

##### **Reducing power** [18]:

Substances which have reduction potential react with potassium ferricyanide (Fe<sup>3+</sup>) to form potassium ferrocyanide (Fe<sup>2+</sup>) which then reacts with ferric chloride

to form ferric-ferrous complex that has an absorption maximum at 700 nm. Different doses of Ethanolic extract of BO were mixed in 1 ml of distilled water so as to get 20, 40, 60, 80, 100 ug concentration. This was mixed with phosphate buffer (2.5 ml, 0.2 M, pH 6.6) and potassium ferricyanide (2.5 ml, 1%). The mixture was incubated at 50°C for 20 min. A portion (2.5 ml) of trichloro acetic acid (10%) was added to the mixture, which was then centrifugated at 3000 rpm for 10 min. The upper layer of the solution (2.5 ml) and FeCl<sub>3</sub> (0.5 ml, 0.1%) and the

absorbance was measured at 700 nm. Increased absorbance of the reaction indicates increase in reducing power. Ascorbic acid was used as standard. The % reducing power was calculated by using the formula:

$$\% \text{ increase in the absorbance} = \frac{\text{Control OD} - \text{Test OD}}{\text{Control OD}} \times 100$$

The results are shown in **Table 4** and depicted in **figure 4**


**RESULTS:**

**Table 1: Effect of BO Extract on Immobility Time in Fst in Rats**

Sr. No.	Group	Immobility Time In Seconds
1	Control 1% CMC	205 + 3.2
2	Standard Imipramine (20 mg/kg)	249 + 2.19
3	BO Extract 10mg/kg	199 + 6.4*
4	BO Extract 25mg/kg	211 + 3.3**
5	BO Extract 50mg/kg	232 + 4.0**

The Results are expressed as mean of six animals, n=6

\* P < 0.05 when compared to control \*\* P < 0.001 when compared to control



**Figure 1: Effect of BO Extract on Immobility Time in Fst in Rats**

**Table 2: DPPH radical scavenging activity of BO extract**

Concentration of extract (µg/ml)	% Inhibition			
	AA	BO	AA	BO
0 (Control)	0.884± 0.002	0.876± 0.001	0	0
10	0.728 ± 0.002*	0.795±0.004	30.08	23.4
20	0.517 ± 0.002*	0.602±0.002*	54.06	31.2
40	0.392 ± 0.003*	0.515±0.005*	68.7	55.3
60	0.251 ± 0.005*	0.360±0.003*	84.7	73.0
80	0.193 ± 0.002*	0.304±0.001*	91.2	79.4
100	0.140 ± 0.005*	0.272±0.003*	97.2	83.1

Values are Mean of three replicates. Statistical analysis by ANOVA followed by Dunnett multiple comparison test. Significance at \*p<0.01 when compared to control.




Figure 2: DPPH radical scavenging activity of BO extract

Table 3: Nitric oxide scavenging method of BO extract

Concentration of extract (ug/ml)	% Inhibition			
	AA	BO	AA	BO
0 (Control)	0.869±0.008	0.865±0.004	0	0
20	0.646 ±0.003*	0.735±0.002*	40.7	30.0
40	0.436 + 0.003*	0.530±0.005*	64.9	54.3
60	0.252 ± 0.004*	0.445±0.006*	86.07	64.6
80	0.189 ± 0.007*	0.361±0.002*	93.2	73.8
100	0.147 ±0.003*	0.264±0.008*	98.1	82.2

Values are Mean of three replicates, Statistical analysis by ANOVA followed by Dunnett multiple comparison test. \*p<0.001 when compared to control.




Figure 3: Nitric Oxide radical scavenging activity of BO extracts

Table 4: Reducing Power method of PV and BO extracts

Concentration of extract (ug/ml )	% Inhibition			
	AA	BO	AA	BO
0 (Control)	0.852±0.004	0.852±0.004	0	0
10	0.041±0.005*	0.078±0.003*	95.1	90.8
20	0.052±0.001*	0.096±0.003*	93.8	89.4
40	0.120±0.004*	0.165±0.004*	85.9	80.6
60	0.194±0.008*	0.324±0.007*	77.2	61.9
80	0.372±0.002*	0.512±0.001*	56.3	39.9
100	0.506±0.001*	0.638 ±0.008*	40.6	25.1

Values are Mean of three replicates Statistical analysis by ANOVA followed by Dunnett multiple comparison test.\*p < 0.01 when compared to control.




Figure 4: Reducing power of BO extract

Animals were observed for mortality by acute toxicity test in mice for 48 hrs. No mortality was observed and the BO extract was found to be safe up to a dose of 2000 mg/kg. BO extract result decrease in immobility time in concentration dependent manner, the extract at 50 mg/kg shows significant decrease in immobility ( $p < 0.001$ ) as compared to control (Table 1 and Figure 1). The results suggest that the BO extract posses anti-stress activity. It may be due to the presence of phenolic acids in *Borago officinalis* [7] posses immune stimulatory, anti inflammatory effects and used in the treatment of seborrhic dermatitis [8, 9].

In the DPPH radical scavenging assay the drug at various concentrations of ethanolic solution of the drug produced inhibition of DPPH in dose dependent manner. The obtained results of absorbance and % inhibition showed decrease in the concentration of DPPH radical due to scavenging ability of extracts and at 100 µg/ml of ethanolic extract exhibits 83.1% inhibition  $IC_{50}$  values were found to be 41.5 µg/ml. When compared with reference standard Ascorbic acid ( $IC_{50}$  29.9 µg/ml) the scavenging effects were lesser (Table 2 and Figure 2). The nitric oxide generated from sodium nitro prusside when react with oxygen from nitrite, which inhibited by antioxidants by competing with oxygen to react with

nitric oxide. The % inhibition and IC<sub>50</sub> values found to be 82.2% and 42.5 µg/ml.

The BO extract exhibits significant antioxidant activity with low IC<sub>50</sub> values in these two methods. Ascorbic acid which is used as a reference compound showed 50 % inhibition at 35.6 µg/ml (Table 3 and Figure 3). BO extract showed increase in absorbance indicated significant increase in reducing power and results in decrease in percentage inhibition. Compared with ascorbic acid, the reducing power of BO extract was lesser (Table 4 figure 4).

#### DISCUSSION:

There is extensive evidence that single-dose administration of adaptogens activates corticosteroid formation, and that repeated dosage with adaptogens normalizes the levels of stress hormones, such as adrenocorticotrophic hormone (ACTH). It is known that the blood level of corticosteroids increases as a result of long-term training or adaptation, and that a trained organism responds to stress stimuli with only mildly increased activity of the hypothalamic-pituitary axis (HPA), as opposed to a very pronounced increase in activity seen in untrained states. In other words, adaptogens apparently increase the ability of the stress system to respond to stress stimuli in a manner that tends to preserve homeostasis, particularly by modulating the biosynthesis of eicosanoids - including prostaglandins E2 and F2, 5-hydroxyeicosatet racenoic acid (5-HETE), 12-HETE, and leukotriene B4. Moreover, adaptogens also appear to regulate the basal level of the arachidonic acid and also to do this under various stressful conditions, such as immobilization, heavy physical exercise, and radiation injury. Although there is a difference in the mode of action and pharmacologic activity of different adaptogens.

The primary site of action of adaptogens appears to be the HPA, and their secondary sites of action the liver and components of the immune and cardiovascular systems. The excessive activity of the stress system is associated with increased arousal or anxiety, increased blood pressure, gastrointestinal dysfunction and suppression of the immune response. Adaptogens are also thought to function primarily due to their antioxidant and free radical scavenging effects<sup>[1, 2]</sup>. Free radicals are chemical entities that can exist separately with one or more unpaired electrons. Antioxidants may offer resistance oxidative stress by scavenging free radicals. Free radical or reactive oxygen species (ROS) are produced during biochemical redox reactions as part of normal physiological cell metabolism (protection from infectious organism) and as a response to environmental factors

such as UV light, cigarette smoke, environmental pollutants and gamma radiations. Once formed, ROS attack cellular components causing damage to lipids, proteins and DNA, which can initiate numerous diseases, including cancer, atherosclerosis, rheumatoid arthritis, diabetes, liver damage and central system disorders. Living organisms have a large number of antioxidants, including macro and micro molecules and enzymes which represent total antioxidant activity of the system plays a central role in preventing oxidative stress<sup>[3, 4]</sup>.

The results suggest that the BO extract possesses anti-stress activity. It may be due to the presence of phenolic acids *Borago officinalis*<sup>[7]</sup> possesses immune stimulatory, anti-inflammatory effects, anti lymphoma activity<sup>[8, 9]</sup>. The ethanolic extract of BO produced significant % inhibition in the levels of DPPH and NO and significant increase in reducing power. In conclusion, the present study shows that BO extract possesses adaptogenic activity and antioxidant activity. Further studies are in progress to identify the mechanisms underlying the pharmacological activity observed.

#### ACKNOWLEDGEMENT:

The authors are thankful to management of Krupanidhi College of Pharmacy for providing the facilities for the research work.

#### REFERENCES:

1. Winston, David and Maimes, Steven. "Adaptogens: Herbs for Strength, Stamina and Stress Relief," Healing Arts Press, 2007; 24:187-191.
2. Robyn Klein."Allostasis Theory and Adaptogenic Plant Remedies." 2004; 35: 165-172.
3. Marx JL, Oxygen free radicals linked to many diseases. Journal of Plant products. 1987; 235: 529 - 534.
4. The Unani Pharmacopeia of India. published by Department of AYUSH, Ministry of Health & Welfare. New Delhi. 2009; Part-1, 2: 35-36.
5. [http:// earthnotes. tripod. Com / borage \\_ h. htm.](http://earthnotes.tripod.com/borage_h.htm) (Accessed on 21 March 2012)
6. Kirtikar KR, Basu, BD. *Borago officinalis* In: Indian Medicinal Plants. 1980; 3:1672-1702.
7. Mahinda W, Fereidoon S. Anti oxidant and free radical scavenging properties of Ethanolic extracts of defatted borage seeds. Food Chem. 1999; 67(4): 399-414.
8. Van Gool CJ, Thijs C, Henquet CJ et al. Gamma-linolenic acid supplementation for prophylaxis of atopic dermatitis-a randomized controlled trial in infants at high familial risk. American Journal of Clinical Nutrition. 2003; 77(4): 943-951.

9. Ziboh VA, Naguwa S, Vang K et al. Suppression of leukotriene B4 generation by ex-vivo neutrophils isolated from asthma patients on dietary supplementation with gammalinolenic acid-containing borage oil: possible implication in asthma. *Clinical and Developmental Immunology*. 2004; 11(1): 13-21.
10. Organization of Economic cooperation and Development (OECD) guidelines for the testing of chemicals. Available from <http://www.oecd.org>
11. Srinivas N, Rajvanu K, Shishruith K, Krishna A. Adaptogenic and nootropic activity of aqueous extract of *Vitis vinifera* (grape seed). An experimental study in rat model. *BMC complementary Alternative Medicine* 2005; 5: 1-8
12. Shaik A, Amal Kumar H, Suresh RN. Evaluation of adaptogenic activity profile of herbal preparation, *Indian Journal of Experimental biology*. 2006; 44: 574 - 579.
13. Laxmi Narasu N, ShantaKumar SM, Shalam MD. Pharmacological and biochemical evidence of the antidepressant effect of the herbal preparation *Trans – 01. Indian journal of experimental biology*. 2007; 39 (5), 231 – 234
14. Porsolt RD, Anton G, Blavet N, Jalire M. Behavioural despair in rats: A new model sensitive to antidepressant treatments. *Eur J Pharmacol* 1978; 47: 379-91
15. Baskar B, Rajeswari V, Satish Kumar T, Invitro antioxidant studies in leaves of *Annona* species. *Ind J of Exp Bio*. 2007; 45: 480 - 485.
16. Kenjale RD, Shah RK, Sathaye SS. Antistress and antioxidant effects of *chlorophytum borivillianum*, *Ind J Exp Bio*. 2007; 45: 974 - 979.
17. Sharma S K, Gupta K V. Free radical scavenging activity of *Ficus racemosa* roots, *Indian J.Pharm.Educ.Res*, 2007; 41: 394-396.
18. Anuradha CV, Kannapan S, Lakshmi Devi SL. Evaluation of invitro antioxidant activity of Indian bay leaf, *Cinnamomum tamala* (Buch. - Ham.) T.Nees & Ebern using rat brain synaptosomes as model system, *Indian J of Exp Bio*, 2007; 45: 378 – 384.