

Journal of Drug Discovery and Therapeutics

Available Online at www.jddt.in

CODEN: - JDDTBP (Source: - American Chemical Society)

Volume 12, Issue 06; 2024, 116-126

Formulation and Evaluation of Phytosomes of Herbal Plant

Bhingare Bayana Bibhishan¹, Dr. Vidhi Jain², Dr. Vivek Subhash Tarate³¹Research Scholar, Department of Pharmacy, Sabarmati University, Ahmedabad, Gujarat²Research Supervisor, Department of Pharmacy, Sabarmati University, Ahmedabad, Gujarat³Research Co-Supervisor, Department of Pharmacy, Sabarmati University, Ahmedabad, Gujarat

Received: 11-09-2024 / Revised: 11-10-2024 / Accepted: 20-11-2024

Corresponding author: Bhingare Bayana Bibhishan

Conflict of interest: No conflict of interest.

Abstract:

Phyllanthus reticulatus phytosomes were developed and tested in this work with the hope of improving the bioavailability, stability, and therapeutic effectiveness of the plant's bioactive chemicals. After collecting and drying the aerial portions of *P. reticulatus*, a hydroalcoholic solvent was used for extraction. Significant, concentration-dependent activity was observed in the in-vitro antioxidant evaluation of the extract (HEPR) using DPPH and hydroxyl radical scavenging assays. Critical formulation variables such as the phospholipid:cholesterol ratio, temperature, and sonication time were considered in the optimization of HEPR-loaded phytosomes using Box-Behnken Design. The phytosomes were manufactured utilizing the antisolvent precipitation method. The improved phytosomes showed homogeneous dispersion, spherical shape, and nanosized vesicles (157-253 nm) with a high entrapment effectiveness (55-73%). Physical and chemical analyses with DLS, TEM, FTIR, and DSC verified that HEPR was successfully integrated into the phospholipid matrix without causing any chemical reactions. Phytosomal formulation has the ability to enhance antioxidant delivery and therapeutic efficacy, as demonstrated by in-vitro release tests that demonstrated sustained drug release relative to plain extract.

Keywords: Phyllanthus reticulatus, Hydroalcoholic extract (HEPR), Phytosome, Antioxidant activity, Vesicle size, Entrapment efficiency, Box–Behnken design, Controlled drug release

Introduction

Chronic diseases including cancer, heart problems, and dementia are accelerated by oxidative stress, which results from an imbalance between endogenous antioxidants and reactive oxygen species. Because of their bioactive components and lack of adverse effects, medicinal plants show great promise as a natural antioxidant source. The tropical plant *Phyllanthus reticulatus* contains tannins, phenolics, and flavonoids, all of which

have the ability to protect the liver, reduce inflammation, and slow the aging process.

However, due to its fast metabolism, low bioavailability, and poor solubility, *P. reticulatus* extract has limited clinical applicability despite these benefits. Improved absorption, stability, and controlled release of active chemicals are achieved by phytosome technology, which entails complexing phytoconstituents with phospholipids.

This research involved the preparation of HEPR, or hydroalcoholic extract of *P. reticulatus*. After that, HEPR-loaded phytosomes were created, optimized with a Box-Behnken Design, and studied for a variety of properties, including morphology, in-vitro drug release, physicochemical stability, vesicle size, and entrapment efficiency. A new phytosomal delivery system with enhanced antioxidant activity and therapeutic promise was the intended outcome of this strategy.

Materials and method:

For this planned research, we used the aerial portions of *Phyllanthus reticulatus*. The plant parts were gathered from the surrounding region. The plant portions were carefully washed after harvest to eliminate any debris or dirt. After cleaning, the plant material was dried in a shaded area to maintain its phytoconstituents and avoid their deterioration.

The plant material was sun-dried, then pounded into a coarse powder. It was first macerated with petroleum ether to remove any fat. The extraction process was maintained until all lipids were removed from the sample. Then, 30 g of dried *Phyllanthus reticulatus* aerial portions were completely extracted using the maceration process and a hydroalcoholic solvent (ethanol: water, 70:30).

Formulation and Characterization of HEPR Phytosome

Design-Expert software was utilized to optimize the formulation parameters and to investigate the effects of formulation variables on the phytosome characteristics. A Box–Behnken Design (BBD) was employed for the experimental investigation using Design-Expert® trial version 13.0.2.0 (Stat-Ease Inc., Minneapolis, USA).

The Box–Behnken design consisted of three factors at three levels, which were used to optimize the critical material attributes (CMAs) and evaluate their impact on the critical quality attributes (CQAs). The factor levels were coded as -1, 0, and +1, and the corresponding numerical values are presented in Table 3.1.

As shown in the table, the independent variables (CMAs) included the phospholipid:cholesterol ratio (X_1), temperature (X_2), and sonication time (X_3). The dependent variables (CQAs) selected for this study were vesicle size (Y_1) and entrapment efficiency (Y_2).

Direct response values obtained from different experimental runs showing the effect of Amount of Phospholipid: Cholesterol (X_1), Temperature (X_2), and Sonication Time (X_3) on critical quality attributes (CQAs).

Table: Effect of formulation and process variables on HEPR phytosome characteristics

Formulation Code	X_1	X_2	X_3
HEPR1	-1	-1	0
HEPR2	0	0	0
HEPR3	0	-1	1
HEPR4	0	0	0
HEPR5	0	-1	-1
HEPR6	0	1	-1
HEPR7	1	0	1
HEPR8	-1	0	1
HEPR9	0	0	0
HEPR10	-1	1	0
HEPR11	1	0	-1
HEPR12	0	0	0

HEPR13	0	1	1
HEPR14	-1	0	-1
HEPR15	1	-1	0
HEPR16	0	0	0
HEPR17	1	1	0

Characterization of HEPR-Loaded Phytosome

Determination of Entrapment Efficiency

High-performance liquid chromatography (HPLC) was employed to determine the entrapment efficiency (EE), defined as the amount of quercetin successfully incorporated into the phytosomal vesicles. The entrapment efficiency of HEPR-loaded phytosomes was determined using the centrifugation method.

Briefly, the phytosomal dispersion was centrifuged at 12,000 rpm at 4 °C to separate the entrapped drug from the untrapped drug present in the supernatant. The transparent supernatant containing free quercetin was carefully removed. To lyse the phytosomal vesicles, 1 mL of 0.1% Triton X-100 was added to the pellet, followed by appropriate dilution using phosphate-buffered saline (PBS, pH 7.4).

The amount of quercetin present in the HEPR phytosome was quantified using HPLC at 280 nm (Singh *et al.*, 2011). The percentage entrapment efficiency was calculated using the following formula:

$$\%EE = \frac{\text{Amount of drug entrapped}}{\text{Total amount of drug added}} \times 100$$

Physicochemical Characterization of HEPR Phytosome

The particle size, polydispersity index (PDI), and zeta potential (ZP) of HEPR-loaded phytosomes were measured using dynamic light scattering (DLS) on a particle size analyzer (Malvern Zeta Master ZEM 5002, Malvern, UK). The measurements were performed at 25 °C using the laser diffraction method, which detects backscattered laser light.

Zeta potential was determined by measuring the velocity and direction of nanoparticle movement under an applied electric field. The results were expressed as mean \pm standard deviation. All measurements were performed in triplicate ($n = 3$) to ensure accuracy and reproducibility (Anwar *et al.*, 2018; Wan *et al.*, 2019).

Morphology of HEPR Phytosome

The surface morphology and shape of the optimized HEPR phytosome formulation were initially examined using an optical microscope (Cippon, Japan) equipped with a camera attachment (Minolta). A small quantity of the phytosome sample was placed on a glass slide and covered with a coverslip. The sample was observed under 10 \times magnification, and images were captured.

Further morphological evaluation and particle size analysis were conducted using transmission electron microscopy (TEM). A drop of the phytosome dispersion was placed on a carbon-coated copper grid and allowed to stand for 15 minutes. The sample was then negatively stained using 1% aqueous phosphotungstic acid and air-dried. The samples were visualized using a TEM (Hitachi H-7500, Tokyo, Japan) (Yu *et al.*, 2014).

FTIR Studies

(Mazumder *et al.*, 2016; Macedo *et al.*, 2021)

Fourier Transform Infrared (FTIR) spectroscopy was performed to evaluate the compatibility between HEPR and formulation excipients. KBr pellets were prepared by mixing the crude HEPR and optimized HEPR phytosomes with

potassium bromide in a 1:100 ratio using a conventional pellet press technique.

The prepared pellets were placed in the IR chamber, and the spectra were recorded over a wavelength range of 4000–400 cm^{-1} . FTIR spectra were analyzed to identify characteristic functional groups and possible drug–excipient interactions.

DSC Studies

Differential Scanning Calorimetry (DSC) was used to study the thermal behavior of crude HEPR and HEPR-loaded phytosomes. Samples weighing 5–10 mg were sealed in flat-bottomed aluminum pans and heated at a constant rate of 10 $^{\circ}\text{C}/\text{min}$ over a temperature range of 50–400 $^{\circ}\text{C}$ under a nitrogen atmosphere (flow rate: 200 mL/min). Alumina was used as the reference standard (Rasaei *et al.*, 2014; Sohail *et al.*, 2016).

In-Vitro Drug Release Studies

The dialysis membrane technique was employed to study the in-vitro drug release behavior of plain HEPR solution and HEPR-loaded phytosomes. A known quantity of phytosomal formulation (equivalent to 1 mg/mL of HEPR) was placed inside a dialysis membrane, which was then immersed in a glass beaker containing 200 mL of 0.1 N HCl (pH 1.0).

The dissolution medium was maintained at 37 ± 0.5 $^{\circ}\text{C}$ with continuous stirring at 75 rpm. Aliquots of 2 mL were withdrawn at predetermined time intervals (1, 2, 4, 6, 8, 10, and 12 hours) and immediately replaced with fresh dissolution medium to maintain sink conditions.

The withdrawn samples were filtered using Whatman filter paper and analyzed by HPLC at 280 nm to quantify the amount of

quercetin released. All experiments were conducted in triplicate. The drug release data were fitted to various kinetic models, including zero-order, first-order, Higuchi, and Korsmeyer–Peppas models, to determine the release mechanism (Karole *et al.*, 2019).

Results and discussion:


Fabrication and Evaluation of HEPR Phytosome

The antisolvent precipitation method was employed for the fabrication of HEPR phytosomes. The nanosized vesicular system was prepared using a hydroalcoholic extract of *Phyllanthus reticulatus* along with a phospholipid–cholesterol mixture. Phospholipids acted as the carrier, while cholesterol imparted rigidity to the vesicular system. The formulation parameters influencing phytosome characteristics included the phospholipid:cholesterol ratio, temperature, and sonication time. These variables were systematically evaluated using a Box–Behnken Design (BBD) (Marasini *et al.*, 2012).

Box–Behnken Design for Optimization of HEPR Phytosome

Statistical optimization of HEPR phytosomes was performed using response surface methodology. A total of 17 experimental batches were prepared, and the obtained data were used to evaluate the effect of formulation variables on vesicle size and entrapment efficiency. Vesicle size ranged from 157.73 ± 4.15 nm to 253.47 ± 6.63 nm, while entrapment efficiency varied from $55.23 \pm 2.65\%$ to $72.76 \pm 1.77\%$. Regression analysis was carried out to understand the relationship between CMAs and CQAs,

Formulation code	CQAs			CMAs	
	X1	X2	X3	Y1 (Vesiclesize, nm)	Y2 (Entrapment efficiency, %)
HEPR1	-1	-1	0	195.74±8.93	60.93±3.91
HEPR2	0	0	0	183.37±7.92	66.8±2.50
HEPR3	0	-1	1	214.08±4.62	64.6±3.16
HEPR4	0	0	0	200.24±5.0	67.57±1.65
HEPR5	0	-1	-1	188.96±2.91	58.86±3.30
HEPR6	0	1	-1	187.3±4.42	60.4±6.91
HEPR7	1	0	1	274.8±7.82	69.8±1.76
HEPR8	-1	0	1	219.46±3.72	67.22±3.70
HEPR9	0	0	0	172.03±5.95	69.22±2.92
HEPR10	-1	1	0	157.73±4.15	62.49±2.50
HEPR11	1	0	-1	251.14±5.69	58.23±3.28
HEPR12	0	0	0	179.06±3.25	69.88±3.35
HEPR13	0	1	1	253.47±6.63	72.76±1.77
HEPR14	-1	0	-1	213.61±2.89	55.23±2.65
HEPR15	1	-1	0	219.11±3.36	61.13±4.48
HEPR16	0	0	0	180.36±5.32	66.7±1.18
HEPR17	1	1	0	238.92±4.14	72.27±1.50



Assessment of CMA Influence on CQAs


Effect on Vesicle Size

The vesicle size (Y_1) was significantly influenced by formulation variables. The developed polynomial equation indicated that the phospholipid:cholesterol ratio (X_1) and sonication time (X_3) had a positive effect on vesicle size, while temperature (X_2) showed a comparatively lower impact. An increase in lipid concentration and sonication time resulted in larger vesicles due to enhanced vesicle interaction and agglomeration. These effects were clearly depicted in the perturbation, 2D, and 3D response surface


plots (Bouarab *et al.*, 2014; Ahmed *et al.*, 2019).

Effect on Entrapment Efficiency


The entrapment efficiency (Y_2) increased with higher levels of phospholipid concentration and sonication time, as represented by the coded polynomial equation generated using Design-Expert software. Increased phospholipid content enhanced drug-loading capacity, while longer sonication time promoted better vesicle formation and drug encapsulation. Temperature also played a supportive role in improving entrapment efficiency. The interaction effects were visualized through 2D and 3D response plots




Particle size of optimized HEPR Phytosome



A) Surface morphology of HEPR Phytosome using Optical microscope, (B) TEM



FTIR Spectra of HEPR and HEPR Phytosomes




DSC thermogram of HEPR and HEPR Phytosomes

In-vitro dissolution study of phytosomes


Time (hrs)	Sq.rt.of Time	Log Time	%Cumulative drug release of HEPR loaded phytosomes	%Cumulative drug release of HEPR
1	1	0.00	19.47±0.53	12.32±0.37
2	1.41	0.30	29.08±0.68	20.21±0.43
4	2	0.60	42.51±0.32	32.62±0.18
6	2.44	0.77	57.28±0.71	45.87±0.52
8	2.82	0.90	69.71±0.48	53.18±0.84
10	3.16	1.00	81.41±0.27	62.93±0.63
12	3.46	1.07	94.79±0.93	65.23±0.57

Log cumulative percentage drug release from HEPR loaded phytosomes and HEPR


Time (hrs)	Sq.rt.of Time	Log Time	Log cumulative percentage drug release	
			HEPR loaded phytosomes	HEPR
1	1	0.00	1.28	1.09
2	1.41	0.30	1.46	1.30
4	2	0.60	1.62	1.51
6	2.44	0.77	1.75	1.66
8	2.82	0.90	1.84	1.72
10	3.16	1.00	1.91	1.79
12	3.46	1.07	1.97	1.81



In-vitro drug release profile of HEPR loaded phytosomes and HEPR to zero order release



In-vitro drug release profile of HEPR loaded phytosomes and HEPR according to first order release



In-vitro drug release profile of HEPR loaded phytosomes and HEPR according to Higuchi's model

Conclusion:

The present study was undertaken to evaluate the antioxidant potential of *Phyllanthus reticulatus* extract and to develop and characterize a phytosomal formulation (HEPR phytosome) for improved delivery and therapeutic performance. The hydroalcoholic extract of *Phyllanthus reticulatus* demonstrated significant in-vitro antioxidant activity, as evidenced by DPPH and hydroxyl radical scavenging assays. The antioxidant activity increased in a concentration-dependent manner, indicating the presence of bioactive phytoconstituents such as flavonoids and phenolic compounds,

which are known for their free radical scavenging properties.

To enhance the bioavailability and stability of the extract, HEPR phytosomes were successfully formulated using the antisolvent precipitation technique. Optimization using the Box–Behnken design enabled systematic evaluation of the influence of critical material and process variables on vesicle size and entrapment efficiency. The optimized formulations exhibited nanosized vesicles with acceptable polydispersity and high entrapment efficiency, confirming the suitability of phospholipid–cholesterol combinations for phytosome development.

The vesicle size was significantly influenced by the phospholipid:cholesterol ratio and sonication time. Increased lipid concentration and prolonged sonication resulted in larger vesicles due to enhanced vesicle interaction and agglomeration. Entrapment efficiency improved with higher phospholipid content and sonication time, which can be attributed to increased lipid matrix availability and enhanced drug encapsulation capacity. These findings are in agreement with previously reported phytosomal and nanovesicular studies.

Physicochemical characterization confirmed the formation of stable phytosomes with uniform particle size distribution and sufficient surface charge to prevent aggregation. Morphological studies using optical microscopy and TEM revealed spherical and well-dispersed vesicles, supporting the DLS findings. FTIR and DSC analyses confirmed the compatibility of HEPR with excipients and suggested successful incorporation of the extract within the phytosomal system without any chemical interaction.

In-vitro drug release studies demonstrated that HEPR-loaded phytosomes exhibited higher and sustained cumulative drug release compared to plain HEPR extract. The enhanced release profile may be attributed to improved solubility, increased surface area, and controlled release behavior provided by the phospholipid bilayer. Kinetic modeling indicated that the release pattern followed diffusion-controlled mechanisms, supporting the sustained release characteristics of the phytosomal formulation.

Overall, the findings of this study highlight that phytosomal encapsulation significantly improves the antioxidant potential, stability, and release behavior of *Phyllanthus reticulatus* extract. The optimized HEPR phytosome formulation shows promise as an effective natural antioxidant delivery system and may be further explored for in-vivo evaluation and therapeutic applications.

References:

1. Ahmed, S., Kumar, A., & Rizvi, M. M. (2019). Formulation and optimization of phytosomes: A review. *Journal of Drug Delivery Science and Technology*, 52, 753–765.
2. Anwar, S., Iqbal, J., & Rehman, A. (2018). Physicochemical characterization of nanovesicular systems: Particle size, PDI, and zeta potential. *International Journal of Pharmaceutics*, 547(1–2), 1–10.
3. Bouarab, L., Lakhdar, A., & Benajiba, N. (2014). Effect of lipid concentration on vesicle size and stability of phytosomes. *Journal of Nanoscience and Nanotechnology*, 14(7), 524–531.
4. Karole, S., Gupta, A., & Sharma, R. (2019). In-vitro drug release kinetics: Models and applications. *Journal of Pharmaceutical Analysis*, 9(5), 328–336.
5. Marasini, N., Skalko-Basnet, N., & Sefcik, J. (2012). Application of Box–Behnken design in nanoparticle formulation optimization. *International Journal of Nanomedicine*, 7, 2071–2083.
6. Maryana, H., Supriatna, A., & Wibowo, A. (2016). Effect of sonication and lipid content on vesicle size and entrapment efficiency. *Pharmaceutics*, 8(2), 17.
7. Mazumder, U. K., Gupta, M., & Saha, S. (2016). FTIR and DSC studies of drug–phospholipid complexes. *Journal of Pharmaceutical Sciences*, 105(3), 987–995.
8. Rasaei, M. J., Shojaei, M., & Ebrahimi, S. (2014). Thermal and physicochemical characterization of phytosomes. *Journal of Thermal Analysis and Calorimetry*, 117, 1121–1128.
9. Singh, P., Sharma, B., & Kumar, V. (2011). HPLC estimation of flavonoids in phytosome formulations. *Journal of Chromatographic Science*, 49(5), 347–352.

10. Sohail, M., Ahmed, M., & Khan, F. (2016). Differential scanning calorimetry for nanocarrier characterization. *Journal of Thermal Analysis and Calorimetry*, 126, 345–352.
11. Wan, X., Li, Y., & Zhao, Y. (2019). Characterization of nanoscale phytosomes using dynamic light scattering. *Journal of Nanobiotechnology*, 17, 1–12.
12. Yu, J., Liu, X., & Gao, Y. (2014). TEM characterization of drug-loaded nanovesicles. *Micron*, 65, 9–16.
13. Guchu, S. M., Machocho, A. K., Mwihi, J. T., & Ngugi, M. P. (2020). In-vitro antioxidant activities of methanolic plant extracts. *Journal of Phytopharmacology*, 9(2), 85–91.
14. Gupta, M., Mazumder, U. K., Sivakumar, T., & Vamsi, M. L. M. (2007). Antioxidant and anti-inflammatory activities of *Phyllanthus niruri*. *Journal of Ethnopharmacology*, 114, 364–370.
15. Bouayed, J., Rammal, H., & Soulimani, R. (2009). Oxidative stress and antioxidant bioactive compounds: Mechanisms and health benefits. *Current Nutrition & Food Science*, 5(4), 1–15.