

**IMMUNE RESPONSE DRUG CONJUGATES: CARBON-14 LABELING NECESSITIES**

Pramod Singh Khatri

PhD. Scholar, Department of Pediatrics, AIIMS Hospital, India.

ABSTRACT

Despite recent advances in early detection and treatment, prostate cancer is the most common cancer and the second leading cause of cancer-related deaths in developed countries. Prostate cancer patients having a limited number of treatment options such as radiotherapy, cytotoxic chemotherapy etc. Although results of all approaches are showing beneficial effects, but without long term survival benefit. A recent development in the field of cancer treatment named as gene therapy comes in existence with promise of prostate cancer treatment. It is an emerging field against not only prostate cancer but for all types of cancers. With advancement of basic research and gene therapy technology, no doubt it may be the best future cancer treatment with the use of genetic engineering and clinical trials. In this review we tried to conclude the importance of the gene therapy in prostate cancer with novel and unique characteristics and with promise to cure as the hope for management of the prostate cancer.

KEYWORDS: Gene Therapy, Carcinoma Prostate, Gene Transfer, Immunotherapy, Oncolytic Virotherapy.

INTRODUCTION:

Experiments on animals are still necessary for explanation of fundamental mechanism of the malignancy and to find out the improved methods to prevent, diagnose and treat cancer (1). Despite recent advances in early detection and treatment, prostate cancer is the most common cancer and the second leading cause of cancer-related deaths in developed countries (2,3). Prostate cancer patients having a limited number of treatment options such as radiotherapy, cytotoxic chemotherapy etc. Although results of all approaches are showing beneficial effects, but without long term survival benefit (4-6). Unfortunately, with all possible treatments of the prostate cancer, the patients become non-responsive towards treatment with high mortality due to uncontrolled death. Regarding this issue it is necessary to develop and use some new more effective therapeutic approaches. Present treatments often showing long term negative side effects (7). In addition, Chemotherapy causes long-term side effects, including an increased risk of developing other types of cancers (6,8). In case of metastatic prostate cancer treatment, on one hand treatment prolonging life and on the other hand causes negative side effects like: hot flashes, impotence, incontinence, an increased risk of bone fractures etc. (9). Therefore, it is a need for new treatment option to reduce the death due to prostate cancer.

One new therapy named as gene therapy comes in existence with promise of prostate cancer treatment (10-12). The emerging field of cancer treatment "gene therapy" offers a number of exciting potential treatments (13). Wide range of treatment options occurs with use of genetic material either in vitro or in vivo to cure the prostate

cancer in gene therapy (13). Under this therapeutic approach potential therapeutic gene injected directly in to tumor. These genes directly act on the target genes resulting monitored disease progression. Recently, several gene therapy strategies have generated provocative results in early-stage clinical trials, raising the possibility that gene therapy may have the potential to affect both localized and metastatic disease (14). This disease progression monitored by measuring prostate-specific antigen (PSA) (10-12,15-16). Prostate cancer is amenable in nature, relatively slow-growing disease. With a plus point, due to amenable and slow-growing nature, multiple gene therapies can be used to for the prostate cancer treatment (17-19).

In this review we tried to conclude the importance of the gene therapy in prostate cancer with novel and unique characteristics and with promise to cure as the hope for management of the prostate cancer.

GENE THERAPY:

Lack of significant efficacy is a major limitation of human cancer gene therapy (20). Gene therapy is a broad field with a dedicated promise for cure and prevention of deaths due to prostate cancer. Gene therapy comprises by means three types of techniques; Immunotherapy, Oncolytic Virotherapy and Gene Transfer. These all techniques are used independently for the treatment of the prostate cancer. We hope, with more research in the field of gene therapy, all the techniques may be used in combination for treatment of not even the prostate cancer but all type of prostate cancers (21). We cannot ignore the importance of the Adenovirus in the gene therapy because

these are the most common vehicle for the gene therapy approaches (10,12,22). In recent time, in search of novel and safer prostate cancer treatment, gene therapy delivery agents have been created and a number of prostate cancer patients globally have participated in gene therapy trials with remarkably few treatment side effects (23,24). The most frequent side effects are fever and symptoms that resemble a cold. If the agent is injected, there is often localized swelling and inflammation at the site of the injection (23, 25-27). If we see the comparative results of side effects, other treatments and treatment benefits by gene therapy, side effects are less than the benefits. So we can recommend gene therapy as a conventional treatment (21). Three major issues i.e. systemic delivery, specific introduction, and specific expression of the target gene are to be focused to develop a clinically relevant treatment strategy (20).

With reference to the Genetic Modification Clinical Research Information System (<http://www4.od.nih.gov/oba/RAC/GeMCRIS/GeMCRIS.htm>), more than about 70 gene therapy protocols have been reviewed by the Recombinant DNA Advisory Committee of the National Institutes of Health, targeting with the prostate cancer till 2007. It is not a surprise regarding few proceeded towards the clinic and even one or two have shown results (14). Question arises regarding what is the main cause of this failure? The answer is very logical i.e. because of lack of resources, regulatory barriers, etc. (28). Both approaches in vivo as well as ex vivo, regarding the gene therapy trials reported till date for the prostate cancer treatment may be an attractive target for the further research. This is because of the various strategies required to promise for localized and metastatic disease. This discussion regarding the localized versus metastatic prostate cancer treatment strategies can be discussed separately.

GENE THERAPY FOR LOCALIZED PROSTATE CANCER:

Gene therapy for localized prostate cancer initiated from replication-defective adenoviruses containing a single therapeutic gene to replication-competent oncolytic adenoviruses containing multiple therapeutic genes (29-37). Freytag and colleagues in 2007 evaluated in the setting of locally recurrent and newly diagnosed disease, and as single agents or in combination with radiation therapy (14). In all, low toxicity in humans has demonstrated. Freytag et al. were the first to “arm” a replication-competent adenovirus with a therapeutic gene and propose combining gene therapy and oncolytic viral therapy with radiotherapy (38-41). In a study by the Freytag, primary results increase the possibility that replication-competent adenovirus-mediated suicide gene therapy may have the potential to

improve local tumor control of conformal radiotherapy in select patient groups (42).

GENE THERAPY FOR METASTATIC PROSTATE CANCER:

Previously both in vivo and ex vivo gene therapy strategies targeting metastatic prostate cancer have been evaluated in the clinic. Kubo and colleagues were among the first to evaluate an in vivo adenovirus-mediated gene therapy strategy targeting metastatic prostate cancer (43). Simon and colleagues were among the first to evaluate an ex vivo gene therapy strategy targeting metastatic prostate cancer (44). Their strategy utilized an autologous granulocyte–monocyte colony stimulating factor –secreting cancer cell vaccine to treat men who were found to have metastatic prostate cancer at the time of surgery. The rationale for using granulocyte–monocyte colony stimulating factor as a cancer therapy is well founded scientifically and stems from the fact that it has demonstrated the greatest ability to induce durable tumoricidal anti-tumor immune responses in preclinical models (45) Although not a study endpoint, all subjects demonstrated significant declines in serum PSA, which is expected after prostate resection. However, all subjects exhibited disease progression (14). The main factor for the prostate cancer bone metastasis is epithelial and stromal interactions (46). After the metastasis to bone prostate cancer cells begin to express bone specific proteins such as OC, osteopontin, and bone sialoprotein (47). All above gene expressions are controlled by the many factors. These properties are known as osteomimetic properties, which can be a golden opportunity to target bony metastases of prostate cancer (48-54).

IMMUNOTHERAPY:

In this type of gene therapy, genetically modified cells and viral particles are usually used for immune system stimulation resulting cancer cells destruction (21). The main purpose of the immunotherapy is to boost the immune system targeting and destruction of cancer cells (55). Currently there are many new researches going on in the field of gene therapy to create recombinant cancer vaccines (56). It is favorite modality, has proved itself better therapeutically under clinical investigations (12,57). It is described under three common approaches (i) tumor cell vaccines, (ii) vector-based vaccines, and (iii) dendritic cell (DC)-based vaccines (20).

Vaccine models showing positive results for prostate cancer treatment in preclinical trials. For example: in a prostate cancer vaccine trial, a patient who achieved normal prostate specific antigen (PSA) levels for the year of the trial, developed rising PSA levels after vaccination was stopped. His PSA levels were stabilized again only with re-

initiation of the vaccine therapy. (58-59). These results, while not entirely positive, have given scientists a better understanding of the immune reaction to cancer and have led to the development of the next generation of cancer vaccines (21).

ONCOLYTIC VIROTHERAPY:

It is an emerging treatment approach that uses replication-competent viruses to destroy cancers. (60) Under oncolytic virotherapy, viral particles are used for replication inside the cancer cell resulting cell death. This is an emerging treatment modality with great promise for treatment of the metastatic prostate cancer (21). As with many current treatment regimens for cancer, gene therapy utilized as a monotherapy may not be the sole solution. A multimodal approach combining virus based gene therapy with chemotherapy and/or radiotherapy may be necessary for more complete tumor eradication (61). Although the majority of our work to date has utilized the prostate cancer model, researchers have also demonstrated the utility of gene therapy in vitro and/or in vivo for several other tumor models, including breast, pancreas, colon, medullary thyroid, ovarian carcinomas (62-63), and in multiple myeloma (64).

GENE TRANSFER:

This is a recent treatment modality in which new genes introduction takes place into a cancerous cell to cause the cell death or slow down the cancer cells growth. It is a very flexible cancer treatment technique with the help of wide range of genes and vectors used for positive results against the prostate cancer treatment (21). Gene therapy for treatment of cancer is generally substratified into four strategies based on the molecular target of gene transfer (65):

1. Tumor suppressor gene therapy
2. Suicide gene therapy
3. Immune modulatory gene therapy
4. Antioncogene therapy

ROLE IN PROSTATE CANCER TREATMENT:

In recent days there is gradual increase in incidence and mortality of prostate cancer (66). Management of localized prostate cancer is possible with surgery or radiation (67-69). Unfortunately, these treatment types are less effective against more aggressive forms of the disease (Stage XT2b or Gleason score X7 or PSA 410 ng/mL), and many of these patients relapse. Because prostate cancer is usually hormone-dependent at this point, patients with a rising prostate-specific antigen (PSA) following definitive therapy are typically treated with salvage androgen suppression therapy (AST).

Despite of these some therapies challenging sure treatment of the prostate cancer, there is no effective curable approach for prostate cancer in an androgen independent state or aggressive state (70-71). To overcome the present problem of prostate cancer it is a very urgent need to search, invent and develop new and effective therapies (71). With respect to all the therapies present for the prostate cancer treatment, gene therapy is seems to be a ray of therapeutic hope for the prostate cancer treatment (71-72).

CONCLUSION:

Gene therapy is emerging field against not only prostate cancer but for all types of cancers. By the advancement of basic research and gene therapy technology, no doubt it may be the best future cancer treatment with the use of genetic engineering and clinical trials. But many hurdles have to be crossed to reach towards the effective gene therapy. Gene therapy can be the best in future treatment of the cancer with few concerns regarding the safety and benefits.

ACKNOWLEDGMENT:

The studies described in this review were supported by Indian Council of Medical Research (ICMR), Delhi, India.

REFERENCES:

1. Workman P, Aboagye EO, Balkwill F, Balmain A, Bruder G, Chaplin DJ, Double JA, Everitt J, Farningham DAH, Glennie MJ, Kelland LR, Robinson V, Stratford IJ, Tozer GM, Watson S, Wedge SR, Eccles SA (2010) Guidelines for the welfare and use of animals in cancer research. *British Journal of Cancer* 102:1555 – 1577.
2. Damber JE, Aus G (2008) Prostate cancer. *Lancet* 371:1710-1721.
3. Dash Rupesh, Azab Belal, Shen Xue-Ning, Sokhi Upneet K, Sarkar Siddik, Su Zhao-zhong, Wang Xiang-Yang, Claudio Pier Paolo, Dent Paul, Dmitriev Igor P, Curiel David T, Grant Steven, Sarkar Devanand, Fisher Paul B (2011) Developing an Effective Gene Therapy for Prostate Cancer: New Technologies with Potential to Translate from the Laboratory into the Clinic. *Discov Med* 11(56):46-56.
4. DiLorenzo G, DePlacido S (2006) Hormone refractory prostate cancer (HRPC): present and future approaches of therapy. *Int J Immunopathol Pharmacol* 19(1):11-34.
5. Sternberg CN (2002) Highlights of contemporary issues in the medical management of prostate cancer. *Crit Rev Oncol Hematol* 43(2):105-121.
6. Dash Rupesh, Azab Belal, Shen Xue-Ning, Sokhi Upneet K, Sarkar Siddik, Su Zhao-zhong, Wang Xiang-

- Yang, Claudio Pier Paolo, Dent Paul, Dmitriev Igor P, Curiel David T, Grant Steven, Sarkar Devanand, Fisher Paul B (2011) Developing an Effective Gene Therapy for Prostate Cancer: New Technologies with Potential to Translate from the Laboratory into the Clinic. *Discov Med* 11(56):46-56.
7. Chemotherapy and you: A guide to self-help during cancer treatment. National Institutes of Health Web site. Available at: <http://www.cancer.gov/PDF/b21d0a74-b477-41ec-bdc0-a60bbe527786/chemoandyou.pdf>. Accessed July 31, 2006.
 8. Bassal M, Mertens AC, Taylor L, Neglia JP, Greffe BS, Hammond S, Ronckers CM, Friedman DL, Stovall M, Yasui YY, Robison LL, Meadows AT, Kadan-Lottick NS (2006) Risk of selected subsequent carcinomas in survivors of childhood cancer: a report from the Childhood Cancer Survivor Study. *J Clin Oncol* 24:476-483.
 9. Sharifi N, Gulley JL, Dahut WL (2005) Androgen deprivation therapy for prostate cancer. *JAMA* 294:238-244.
 10. Anderson WF (1998) Human gene therapy. *Nature* 392(6679):25-30.
 11. Gopalkrishnan RV, Kang DC, Fisher PB (2001) Molecular markers and determinants of prostate cancer metastasis. *J Cell Physiol* 189(3):245-256.
 12. Mabweesh NJ, Zhong H, Simons JW (2002) Gene therapy of prostate cancer: current and future directions. *Endocr Relat Cancer* 9(2):115-139.
 13. Mulligan RC (1993) The basic science of gene therapy. *Science* 260:926-932.
 14. Freytag Svend O, Stricker Hans, Movsas Benjamin, Kim Jae Ho (2007c) Prostate Cancer Gene Therapy Clinical Trials. *Molecular Therapy* 15(6):1042-1052.
 15. Wu L, Matherly J, Smallwood A, Adams JY, Billick E, Beldegrun A et al. (2001) Chimeric PSA enhancers exhibit augmented activity in prostate cancer gene therapy vectors. *Gene Ther* 8: 1416-1426.
 16. Zhang L, Adams JY, Billick E, Ilagan R, Iyer M, Le K et al. (2002) Molecular engineering of a two-step transcription amplification (TSTA) system for transgene delivery in prostate cancer. *Mol Ther* 5: 223-232.
 17. National Cancer Institute. The nation's investment in cancer research: A plan and budget proposal for fiscal year 2007. Washington, DC: US National Institutes of Health, October 2005. NIH Publication No. 06-5856.
 18. Cancer Facts and Figures 2006. American Cancer Society Web site. Available at: <http://www.cancer.org/downloads/STT/CAFF2006PWSecured.pdf>. Accessed July 31, 2006.
 19. Edwards BK, Brown ML, Wingo PA, Howe HL, Ward E, Ries LA, Schrag D, Jamison PM, Jemal A, Wu XC, Friedman C, Harlan L, Warren J, Anderson RN, Pickle LW (2005) Annual report to the nation on the status of cancer, 1975-2002, featuring population-based trends in cancer treatment. *J Natl Cancer Inst* 97:1407-1427.
 20. Figueiredo Marxa L, Kao Chinghai, Wu Lily (2007) Advances in Preclinical Investigation of Prostate Cancer Gene Therapy. *Molecular Therapy* 15(6):1053-1064.
 21. Cross Deanna, Burmester James K (2006) Gene Therapy for Cancer Treatment: Past, Present and Future. *Clinical Medicine & Research* 4(3):218-227.
 22. Haviv YS, Blackwell JL, Kanerva A, Nagi P, Krasnykh V, Dmitriev I, Wang M, Naito S, Lei X, Hemminki A, Carey D, Curiel DT (2002) Adenoviral gene therapy for renal cancer requires retargeting to alternative cellular receptors. *Cancer Res* 62(15):4273-4281.
 23. Reid T, Warren R, Kirn D (2002) Intravascular adenoviral agents in cancer patients: lessons from clinical trials. *Cancer Gene Ther* 9:979-986.
 24. Peng Z (2005) Current status of gendicine in China: recombinant human Ad-p53 agent for treatment of cancers. *Hum Gene Ther* 16:1016-1027.
 25. Liu M, Acres B, Balloul JM, Bizouarne N, Paul S, Slos P, Squiban P (2004) Gene-based vaccines and immunotherapeutics. *Proc Natl Acad Sci U S A* 101:14567-14571.
 26. Malaeb BS, Gardner TA, Margulis V, Yang L, Gillenwater JY, Chung LW, Macik G, Koeneman KS (2005) Elevated activated partial thromboplastin time during administration of first-generation adenoviral vectors for gene therapy for prostate cancer: identification of lupus anticoagulants. *Urology* 66:830-834.
 27. Pulkkanen KJ, Yla-Herttua S (2005) Gene therapy for malignant glioma: current clinical status. *Mol Ther* 12:585-598.
 28. American Cancer Society (2007). Cancer Facts and Figures 2007. American Cancer Society.
 29. Herman, JR, Adler, HL, Aguilar-Cordova, E, Rojas-Martinez, A, Woo, S, Timme, TL et al. (1999). In situ gene therapy for adenocarcinoma of the prostate: a phase I clinical trial. *Hum Gene Ther* 10: 1239-1249.
 30. Miles, BJ, Shalev, M, Aguilar-Cordova, E, Timme, TL, Lee, HM, Yang, G et al. (2001). Prostate-specific antigen response and systemic T cell activation after in situ gene therapy in prostate cancer patients failing radiotherapy. *Hum Gene Ther* 12: 1955-1967.
 31. Freytag, S, Movsas, B, Aref, I, Stricker, H, Peabody, J, Pegg, J et al. (2007d). Phase I trial of replication-competent adenovirus-mediated suicide gene therapy in combination with IMRT for the treatment of newly-

- diagnosed prostate cancer. *Mol Ther* published online 20 March 2007 (doi: 10.1038/mt.sj.6300120).
32. Teh, BS, Aguilar-Cordova, E, Kernen, K, Chou, CC, Shalev, M, Vlachaki, MT et al. (2001). Phase I/II trial evaluating combined radiotherapy and in situ gene therapy with or without hormonal therapy in the treatment of prostate cancer—a preliminary report. *Int J Radiat Oncol Biol Phys* **51**: 605–613.
 33. DeWeese, TL, van der Poel, H, Li, S, Mikhak, B, Drew, R, Goemann, M et al. (2001). A Phase I trial of CV706, a replication-competent, PSA selective oncolytic adenovirus, for the treatment of locally recurrent prostate cancer following radiation therapy. *Cancer Res* **61**: 7464–7472.
 34. Freytag, SO, Khil, M, Stricker, H, Peabody, J, Menon, M, DePeralta-Venturina, M et al. (2002). Phase I study of replication-competent adenovirus-mediated double suicide gene therapy for the treatment of locally recurrent prostate cancer. *Cancer Res* **62**: 4968–4976.
 35. Freytag, SO, Paielli, D, Wing, M, Rogulski, K, Brown, S, Kolozsvary, A et al. (2002). Efficacy and toxicity of replication-competent adenovirus-mediated double suicide gene therapy in combination with radiation therapy in an orthotopic mouse prostate cancer model. *Int J Radiat Oncol Biol Phys* **54**: 873–885.
 36. Freytag, SO, Stricker, H, Pegg, J, Paielli, D, Pradhan, DG, Peabody, J et al. (2003). Phase I study of replication-competent adenovirus-mediated double suicide gene therapy in combination with conventional dose three-dimensional conformal radiation therapy for the treatment of newly diagnosed, intermediate- to high-risk prostate cancer. *Cancer Res* **63**: 7497–7506.
 37. Teh, BS, Ayala, G, Aguilar, L, Mai, WY, Timme, TL, Vlachaki, MT et al. (2004). Phase I-II trial evaluating combined intensity-modulated radiotherapy and in situ gene therapy with or without hormonal therapy in treatment of prostate cancer-interim report on PSA response and biopsy data. *Int J Radiat Oncol Biol Phys* **58**: 1520–1529.
 38. Kim, JH, Kim, SH, Brown, SL and Freytag, SO (1994). Selective enhancement by an antiviral agent of the radiation-induced cell killing of human glioma cells transduced with HSV-tk gene. *Cancer Res* **54**: 6053–6056.
 39. Khil, MS, Kim, JH, Mullen, CA, Kim, SH and Freytag, SO (1995). Radiosensitization by 5-fluorocytosine of human colorectal carcinoma cells in culture transduced with cytosine deaminase gene. *Clin Cancer Res* **2**: 53–57.
 40. Freytag, SO, Rogulski, KR, Paielli, DL, Gilbert, JD and Kim, JH (1998). A novel three-pronged approach to selectively kill cancer cells: concomitant viral, double suicide gene, and radiotherapy. *Hum Gene Ther* **9**: 1323–1333.
 41. Rogulski, KR, Freytag, SO, Zhang, K, Gilbert, JD, Paielli, DL, Kim, JH et al. (2000). In vivo anti-tumor activity of ONYX-015 is influenced by p53 status and is augmented by radiotherapy. *Cancer Res* **60**: 1193–1196.
 42. Rauen, KA, Sudilovsky, D, Le, JL, Chew, KL, Hann, B, Weinberg, V et al. (2002). Expression of the coxsackie adenovirus receptor in normal prostate and in primary and metastatic prostate carcinoma: potential relevance to gene therapy. *Cancer Res* **62**: 3812–3818.
 43. Kubo, H, Gardner, TA, Wada, Y, Koeneman, KS and Gotoh, A (2003). Phase I dose escalation clinical trial of adenovirus vector carrying osteocalcin promoter-driven herpes simplex virus thymidine kinase in localized and metastatic hormone-refractory prostate cancer. *Hum Gene Ther* **14**: 227–241.
 44. Simons, JW, Mikhak, B, Chang, JF, DeMarzo, A, Carducci, MA, Lim, M et al. (1999). Induction of immunity to prostate cancer antigens: results of a clinical trial of vaccination with irradiated autologous prostate tumor cells engineered to secrete granulocyte-macrophage colony-stimulating factor using ex vivo gene transfer. *Cancer Res* **59**: 5160–5168.
 45. Dranoff, G, Jaffee, E, Lazenby, A, Golumbek, P, Levitsky, H, Brose, K et al. (1993). Vaccination with irradiated tumor cells engineered to secrete murine granulocyte-macrophage colony-stimulating factor stimulates potent, specific, and long-lasting anti-tumor immunity. *Proc Natl Acad Sci USA* **90**: 3539–3543.
 46. Chung LW, Baseman A, Assikis V, Zhau HE (2005) Molecular insights into prostate cancer progression: the missing link of tumor microenvironment. *J Urol* **173**:10–20.
 47. Koeneman KS, Yeung F, Chung LW (1999) Osteomimetic properties of prostate cancer cells: a hypothesis supporting the predilection of prostate cancer metastasis and growth in the bone environment. *Prostate* **39**:246–261.
 48. Gleave M, Hsieh JT, Gao CA, von Eschenbach AC, Chung LW (1991) Acceleration of human prostate cancer growth in vivo by factors produced by prostate and bone fibroblasts. *Cancer Res* **51**:3753–3761.
 49. Festuccia C, Bologna M, Gravina GL, Guerra F, Angelucci A, Villanova I et al. (1999) Osteoblast conditioned media contain TGF- β 1 and modulate the migration of prostate tumor cells and their interactions with extracellular matrix components. *Int J Cancer* **81**:395–403.
 50. Wu HC, Hsieh JT, Gleave ME, Brown NM, Pathak S, Chung LW (1994) Derivation of androgen-independent

- human LNCaP prostatic cancer cell sublines: role of bone stromal cells. *Int J Cancer* 57:406–412.
51. Thalmann GN, Anezinis PE, Chang SM, Zhou HE, Kim EE, Hopwood VL et al. (1994) Androgen-independent cancer progression and bone metastasis in the LNCaP model of human prostate cancer. *Cancer Res* 54:2577–2581.
 52. Chung LW (1995) The role of stromal-epithelial interaction in normal and malignant growth. *Cancer Surv* 23:33–42.
 53. Rhee HW, Zhou HE, Pathak S, Multani AS, Pennanen S, Visakorpi T et al. (2001) Permanent phenotypic and genotypic changes of prostate cancer cells cultured in a three-dimensional rotating-wall vessel. *In Vitro Cell Dev Biol Anim* 37:127–140.
 54. Koenenman KS, Kao C, Ko SC, Yang L, Wada Y, Kallmes DF et al. (2000) Osteocalcin-directed gene therapy for prostate-cancer bone metastasis. *World J Urol* 18:102–110.
 55. Armstrong AC, Eaton D, Ewing JC (2001) Science, medicine, and the future: Cellular immunotherapy for cancer. *BMJ* 323:1289-1293.
 56. Kowalczyk DW, Wysocki PJ, Mackiewicz A (2003) Cancer immunotherapy using cells modified with cytokine genes. *Acta Biochim Pol* 50:613-624.
 57. Tarassoff, CP, Arlen, PM and Gulley, JL (2006) Therapeutic vaccines for prostate cancer. *Oncologist* 11: 451–462.
 58. Emens LA, Jaffee EM (2005) Leveraging the activity of tumor vaccines with cytotoxic chemotherapy. *Cancer Res* 65:8059-8064.
 59. Greiner JW, Zeytin H, Anver MR, Schlom J (2002) Vaccine-based therapy directed against carcinoembryonic antigen demonstrates antitumor activity on spontaneous intestinal tumors in the absence of autoimmunity. *Cancer Res* 62:6944-6951.
 60. Russell Stephen J, Peng Kah-Whye, Bell John C (2012) Oncolytic virotherapy. *Nature Biotechnology* 30, 658–670.
 61. Ahmed KA, Davis BJ, Wilson TM, Wiseman GA, Federspiel MJ, Morris JC (2012) Progress in gene therapy for prostate cancer. *Front Oncol* 2:172.
 62. Scholz IV, Cengic N, Baker CH, Harrington KJ, Maletz K, Bergert ER et al. (2005) Radioiodine therapy of colon cancer following tissue-specific sodium iodide symporter gene transfer. *Gene Ther* 12:272–280.
 63. Dwyer RM, Bergert ER, O' Connor MK, Gendler SJ, Morris JC (2006) Adenovirus-mediated and targeted expression of the sodium iodide symporter (NIS) permits in vivo radioiodide imaging and therapy of pancreatic tumors. *Hum Gene Ther* 17:1–8.
 64. Dingli D, Peng KW, Harvey ME, Greipp PR, O'Connor MK, Cattaneo R, et al. (2004) Image-guided radio virotherapy for multiple myeloma using a recombinant measles virus expressing the thyroidal sodium iodide symporter. *Blood* 103, 1641–1646.
 65. Djavan Bob, Nasu Yasutomo (2001) Prostate Cancer Gene Therapy-What Have We Learned and Where Are We Going? *Rev Urol* 3(4):179–186.
 66. Jemal A, Siegel R, Xu J, Ward E (2010) Cancer statistics, 2010. *CA Cancer J Clin* 60:277-300.
 67. Freytag Svend O, Movsas Benjamin, Aref Ibrahim, Stricker Hans, Peabody James, Pegg Jan, Zhang Yingshu, Barton Kenneth N, Brown Stephen L, Lu Mei, Savera Adnan, Kim Jae Ho (2007a) Phase I Trial of Replication-competent Adenovirus-mediated Suicide Gene Therapy Combined with IMRT for Prostate Cancer. *Molecular Therapy* 15(5):1016–1023.
 68. Freytag Svend O, Stricker Hans, Peabody James, Pegg Jan, Paielli Dell, Movsas Benjamin, Barton Kenneth N, Brown Stephen L, Lu Mei, Kim Jae Ho (2007b) Five-year Follow-up of Trial of Replication competent Adenovirus-mediated Suicide Gene Therapy for Treatment of Prostate Cancer. *Molecular Therapy* 15(3):636–642.
 69. Peschel RE, Colberg JW (2003) Surgery, brachytherapy, and external-beam radiotherapy for early prostate cancer. *Lancet Oncol* 4:233-241.
 70. Antonarakis ES, Carducci MA (2010) Future directions in castrate-resistant prostate cancer therapy. *Clin Genitourin Cancer* 8:37-46.
 71. Chen Yue, Wang Gang, Kong Deling, Zhang Zhihong, Yang Kuo, Liu Ranlu, Zhao Weiming, Xu Yong (2012) In vitro and in vivo double-enhanced suicide gene therapy mediated by generation 5 polyamidoamine dendrimers for PC-3 cell line. *World Journal of Surgical Oncology* 10:3.
 72. Tangney M, Ahmad S, Collins SA, O'Sullivan GC (2010) Gene therapy for prostate cancer. *Postgrad Med* 122:166-180.